


Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1




3702
3703
3704
3705
3706
3707
3708
3709
3710
3711
3712
3713
3714
3715
3716
3717
3718
3719
3720
3721

X. L. Yu Liu. Carbonylation and loss-of-function analyses of SBPase reveal its metabolic interface role in oxidative stress, carbon assimilation, and multiple aspects of growth and development in Arabidopsis. <i>Mol Plant</i> . 2012. 5:1082-99
C. Bartels Dinakar. Light response, oxidative stress management and nucleic acid stability in closely related Linderniaceae species differing in desiccation tolerance. <i>Planta</i> . 2012. 236:541-55
S. Zeng Yang. AtACDO1, an ABC1-like kinase gene, is involved in chlorophyll degradation and the response to photooxidative stress in Arabidopsis. <i>J Exp Bot</i> . 2012. 63:3959-73
H. Chu Chen. Overexpression of AtOGG1, a DNA glycosylase/AP lyase, enhances seed longevity and abiotic stress tolerance in Arabidopsis. <i>J Exp Bot</i> . 2012. 63:4107-21
M. Feng Duan. Overexpression of thylakoidal ascorbate peroxidase shows enhanced resistance to chilling stress in tomato. <i>J Plant Physiol</i> . 2012. 169:867-77
H. Jin Jeon. Chloroplast-targeted bacterial RecA proteins confer tolerance to chloroplast DNA damage by methyl viologen or UV-C radiation in tobacco (<i>Nicotiana tabacum</i>) plants. <i>Physiol Plant</i> . 2013. 147:218-33
S. Kaul Mahanty. Biochemical and molecular analyses of copper-zinc superoxide dismutase from a C4 plant <i>Pennisetum glaucum</i> reveals an adaptive role in response to oxidative stress. <i>Gene</i> . 2012. 505:309-17
R. D. Blanco Ceccoli. Flavodoxin displays dose-dependent effects on photosynthesis and stress tolerance when expressed in transgenic tobacco plants. <i>Planta</i> . 2012. 236:1447-58
A. Ogata Gaber. The involvement of Arabidopsis glutathione peroxidase 8 in the suppression of oxidative damage in the nucleus and cytosol. <i>Plant Cell Physiol</i> . 2012. 53:1596-606
V. Senthil-Kumar Ramegowda. Expression of a finger millet transcription factor, EcNAC1, in tobacco confers abiotic stress-tolerance. <i>PLoS One</i> . 2012. 7:e40397
S. Wang Xu. RNAi knockdown of rice SE5 gene is sensitive to the herbicide methyl viologen by the down-regulation of antioxidant defense. <i>Plant Mol Biol</i> . 2012. 80:219-35
J. Patrit Dechorgnat. Characterization of the Nrt2.6 gene in Arabidopsis thaliana: a link with plant response to biotic and abiotic stress. <i>PLoS One</i> . 2012. 7:e42491
F. Huaxia Yu. GhWRKY15, a member of the WRKY transcription factor family identified from cotton (<i>Gossypium hirsutum</i> L.), is involved in disease resistance and plant development. <i>BMC Plant Biol</i> . 2012. 12:144
C. W. Lee Li. Arabidopsis root-abundant cytosolic methionine sulfoxide reductase B genes MsrB7 and MsrB8 are involved in tolerance to oxidative stress. <i>Plant Cell Physiol</i> . 2012. 53:1707-19
H. Mhatre Tak. Molecular characterization of VvSDIR1 from <i>Vitis vinifera</i> and its functional analysis by heterologous expression in <i>Nicotiana tabacum</i> . <i>Protoplasma</i> . 2013. 250:565-76
H. Zhai Zhang. Overexpression of rice sphingosine-1-phosphate lyase gene OsSPL1 in transgenic tobacco reduces salt and oxidative stress tolerance. <i>J Integr Plant Biol</i> . 2012. 54:652-62
Q. Zhu Jin. Hydrogen gas acts as a novel bioactive molecule in enhancing plant tolerance to paraquat-induced oxidative stress via the modulation of heme oxygenase-1 signalling system. <i>Plant Cell Environ</i> . 2013. 36:956-69
Y. H. Kim Choe. Homologous expression of gamma-glutamylcysteine synthetase increases grain yield and tolerance of transgenic rice plants to environmental stresses. <i>J Plant Physiol</i> . 2013. 170:610-8
X. Wu Luo. Synergistic effects of GhSOD1 and GhCAT1 overexpression in cotton chloroplasts on enhancing tolerance to methyl viologen and salt stresses. <i>PLoS One</i> . 2013. 8:e54002
J. Zhuo Tan. Nitric oxide mediates cold- and dehydration-induced expression of a novel MfHyPRP that confers tolerance to abiotic stress. <i>Physiol Plant</i> . 2013. 149:310-20

Sedoheptulose-1,7-bisphosphatase (SBPase) is a Calvin cycle enzyme and functions in photosynthetic carbon fixation. We
In the present study, three closely related Linderniaceae species which differ in their sensitivity to desiccation are compa
ABC1 (activity of bc1 complex) is a newly discovered atypical kinase in plants. Here, it is reported that an ABC1 protein ki
Reactive oxygen species (ROS) are toxic by-products generated continuously during seed desiccation, storage, and germi
Photosynthesis provides a strong reducing power and a high risk for generation of reactive oxygen species (ROS) particul
The nature and importance of the DNA repair system in the chloroplasts of higher plants under oxidative stress or UV rad
Superoxide dismutases (SODs) form the foremost line of defense against ROS in aerobes. Pennisetum glaucum cDNA libr
Ferredoxins are iron-sulfur proteins involved in various one-electron transfer pathways. Ferredoxin levels decrease unde
A family of eight genes with homology to mammalian glutathione peroxidase (GPX) isoenzymes, designated AtGPX1-AtG
NAC (NAM, ATAF1-2, and CUC2) proteins constitute one of the largest families of plant-specific transcription factors and
Plant heme oxygenase (HO) catalyzes the oxygenation of heme to biliverdin, carbon monoxide (CO), and free iron (Fe(2+
The high affinity nitrate transport system in Arabidopsis thaliana involves one gene and potentially seven genes from the
BACKGROUND: As a large family of regulatory proteins, WRKY transcription factors play essential roles in the processes o
Excess reactive oxygen species (ROS) accumulation under various environmental stresses can damage intracellular polys
The proteins harboring really interesting new gene (RING) finger domains comprise a large family and play key roles in a
Sphingolipids, including sphingosine-1-phosphate (S1P), have been shown to function as signaling mediators to regulate
Hydrogen gas (H2) was recently proposed as a novel antioxidant and signalling molecule in animals. However, the physio
Various environmental stresses induce reactive oxygen species (ROS), causing deleterious effects on plant cells. Glutathio
In plants, CuZn superoxide dismutase (CuZnSOD, EC 1.15.1.1), ascorbate peroxidase (APX, EC 1.11.1.11), and catalase (CA
Hybrid proline-rich proteins (HyPRPs) are cell wall-localized proteins, and are frequently responsive to environmental str

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant

Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1



3722
3723
3724
3725
3726
3727
3728
3729
3730
3731
3732
3733
3734
3735
3736
3737
3738
3739
3740
3741

J. Mu Li. Paraquat Resistant1, a Golgi-localized putative transporter protein, is involved in intracellular transport of paraquat. <i>Plant Physiol.</i> 2013. 162:470-83
X. S. Wang Huang. A basic helix-loop-helix transcription factor, PtrbHLH, of <i>Poncirus trifoliata</i> confers cold tolerance and modulates peroxidase-mediated scavenging of hydrogen peroxide. <i>Plant Physiol.</i> 2013. 162:1178-94
V. Senthil-Kumar Ramegowda. Drought stress acclimation imparts tolerance to <i>Sclerotinia sclerotiorum</i> and <i>Pseudomonas syringae</i> in <i>Nicotiana benthamiana</i> . <i>Int J Mol Sci.</i> 2013. 14:9497-513
W. H. Wang Sun. Reduction of methylviologen-mediated oxidative stress tolerance in antisense transgenic tobacco seedlings through restricted expression of StAPX. <i>J Zhejiang Univ Sci B.</i> 2013. 14:578-85
J. J. Pickup Wargent. Reduction of photosynthetic sensitivity in response to abiotic stress in tomato is mediated by a new generation plant activator. <i>BMC Plant Biol.</i> 2013. 13:108
X. Hu Deng. TaCIPK29, a CBL-interacting protein kinase gene from wheat, confers salt stress tolerance in transgenic tobacco. <i>PLoS One.</i> 2013. 8:e69881
X. Qin Liang. Genes for iron-sulphur cluster assembly are targets of abiotic stress in rice, <i>Oryza sativa</i> . <i>Plant Cell Environ.</i> 2014. 37:780-94
B. Carron De Coninck. Mining the genome of <i>Arabidopsis thaliana</i> as a basis for the identification of novel bioactive peptides involved in oxidative stress tolerance. <i>J Exp Bot.</i> 2013. 64:5297-307
F. Deng Kong. LeCDJ1, a chloroplast DnaJ protein, facilitates heat tolerance in transgenic tomatoes. <i>J Integr Plant Biol.</i> 2014. 56:63-74
H. L. Ma Feng. A novel tomato MYC-type ICE1-like transcription factor, SlICE1a, confers cold, osmotic and salt tolerance in transgenic tobacco. <i>Plant Physiol Biochem.</i> 2013. 73:309-20
A. Perl-Treves Perl. Enhanced oxidative-stress defense in transgenic potato expressing tomato Cu,Zn superoxide dismutases. <i>Theor Appl Genet.</i> 1993. 85:568-76
A. Bonifacio Caverzan. The knockdown of chloroplastic ascorbate peroxidases reveals its regulatory role in the photosynthesis and protection under photo-oxidative stress in rice. <i>Plant Sci.</i> 2014. 214:74-87
R. Yonekura-Sakakibara Nakabayashi. Enhancement of oxidative and drought tolerance in <i>Arabidopsis</i> by overaccumulation of antioxidant flavonoids. <i>Plant J.</i> 2014. 77:367-79
C. Wei Wang. Down-regulation of OsSPX1 causes high sensitivity to cold and oxidative stresses in rice seedlings. <i>PLoS One.</i> 2013. 8:e81849
J. Xia Zhou. RBOH1-dependent H ₂ O ₂ production and subsequent activation of MPK1/2 play an important role in acclimation-induced cross-tolerance in tomato. <i>J Exp Bot.</i> 2014. 65:595-607
W. Hao Shi. The <i>Gossypium hirsutum</i> WRKY gene GhWRKY39-1 promotes pathogen infection defense responses and mediates salt stress tolerance in transgenic <i>Nicotiana benthamiana</i> . <i>Plant Cell Rep.</i> 2014. 33:483-98
A. Keller Egert. Abiotic stress-induced accumulation of raffinose in <i>Arabidopsis</i> leaves is mediated by a single raffinose synthase (RS5, At5g40390). <i>BMC Plant Biol.</i> 2013. 13:218
P. J. Senthilkumar Chen. Transplastomic <i>Nicotiana benthamiana</i> plants expressing multiple defence genes encoding protease inhibitors and chitinase display broad-spectrum resistance against insects, pathogens and abiotic stresses. <i>Plant Biotechnol J.</i> 2014. 12:503-15
A. Kumar Dalal. Alleviation of methyl viologen-mediated oxidative stress by <i>Brassica juncea</i> annexin-3 in transgenic <i>Arabidopsis</i> . <i>Plant Sci.</i> 2014. 219-220:9-18
J. Y. Kim Cha. NADPH-dependent thioredoxin reductase A (NTRA) confers elevated tolerance to oxidative stress and drought. <i>Plant Physiol Biochem.</i> 2014. 80:184-91

Paraquat is one of the most widely used herbicides worldwide. In green plants, paraquat targets the chloroplast by trans
The basic helix-loop-helix (bHLH) transcription factors are involved in a variety of physiological processes. However, plan
Acclimation of plants with an abiotic stress can impart tolerance to some biotic stresses. Such a priming response has not
Ascorbate peroxidases are directly involved in reactive oxygen species (ROS) scavenging by reducing hydrogen peroxide t
BACKGROUND: Yield losses as a result of abiotic stress factors present a significant challenge for the future of global food
Calcineurin B-like protein-interacting protein kinases (CIPKs) have been found to be responsive to abiotic stress. Howeve
Iron-sulphur (Fe-S) cluster assembly occurs in chloroplasts, mitochondria and cytosol, involving dozens of genes in higher
Although evidence has accumulated on the role of plant peptides in the response to external conditions, the number of p
The roles of a tomato (<i>Lycopersicon esculentum</i>) chloroplast-targeted DnaJ protein (LeCDJ1) were investigated using wild
ICE1 (inducer of CBF expression 1), a MYC-type bHLH transcription factor, is an important activator of CBF3/DREB1A for r
The two cDNAs coding for the cytosolic (cyt) and the chloroplast-located (chl) Cu,Zn superoxide dismutases (SODs) of tom
The inactivation of the chloroplast ascorbate peroxidases (chlAPXs) has been thought to limit the efficiency of the water-
The notion that plants use specialized metabolism to protect against environmental stresses needs to be experimentally
Rice SPX domain gene, OsSPX1, plays an important role in the phosphate (Pi) signaling network. Our previous work show
H ₂ O ₂ and mitogen-activated protein kinase (MAPK) cascades play important functions in plant stress responses, but thei
KEY MESSAGE: Our results indicate that overexpression of the GhWRKY39 - 1 gene enhances resistance to pathogen infec
BACKGROUND: The sucrosylgalactoside oligosaccharide raffinose (Raf, Suc-Gal1) accumulates in Arabidopsis leaves in res
Plastid engineering provides several advantages for the next generation of transgenic technology, including the convenie
Plant annexins function as calcium-dependent or -independent phospholipid binding proteins and constitute about 0.1%
NADPH-dependent thioredoxin reductases (NTRs) are key-regulatory enzymes determining the redox state of the thiorec

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant